

* Clarke's research was partially supported by Grant 5-54891, UBC Research Services, and Sun's work was partially supported by National Security Agency under Grant No. MDA904-96-1-0074 and a Research Council Grant from the University of Missouri-Columbia.

Here we present an asymptotic expansion for the expected value of the posterior evaluated at the true value of a parameter. The result can be generalized to give the expected value of the posterior at any other value of the parameter. The novelty is that the expectation is taken over the sample space, not over the parameter space, and is taken with respect to the distribution indexed by the true value of the parameter. The main contribution, aside from the idea of obtaining these expansions, is the technical effort required to demonstrate their feasibility. The formal results were motivated in part by a proposal attributed to I. J. Good, see Kass and Raftery (1995). Good's proposal was that the Bayes factor be used as a test statistic for a frequentist hypothesis test. Our results apply to the

1. Introduction and summary

Key words and phrases: Expected posterior, asymptotics, relative entropy, chi-squared distance, Bayes factor.

Abstract. Let (X_1, \dots, X_n) be independently and identically distributed observations from an exponential family p_θ equipped with a smooth prior density w on a real d -dimensional parameter θ . We give conditions under which the expected value of the posterior density evaluated at the true value of the parameter, θ_0 , admits an asymptotic expansion in terms of the Fisher information $I(\theta_0)$, the prior w , and their first two derivatives. The leading term of the expansion is of the form $n^{d/2} c_1(d, \theta_0)$ and the second order term is of the form $n^{d/2-1} c_2(d, \theta_0, w)$, with an error term that is $o(n^{d/2-1})$. We identify the functions c_1 and c_2 explicitly. A modification of the proof of this expansion gives an analogous result for the expectation of the square of the posterior evaluated at θ_0 . As a consequence we can give a confidence band for the expected posterior, and we suggest a frequentist refinement for Bayesian testing.

(Received April 30, 1996; revised October 6, 1997)

¹Department of Statistics, University of British Columbia,
6356 Agricultural Road, Vancouver, Canada V6T 1Z2
²Department of Statistics, 222 Math Science Building, University of Missouri-Columbia,
Columbia, MO 65211, U.S.A.

BERTRAND CLARKE¹ AND DONGCHU SUN²

ASYMPTOTICS OF THE EXPECTED POSTERIOR*

Ann. Inst. Statist. Math.
Vol. 51, No. 1, 163-185 (1999)

expectation of a posterior density, not to the ratio of two posterior probabilities directly. Nevertheless, our results provide a partial frequentist characterization of the sampling distribution for a Bayesian hypothesis test. Consequently, we give heuristic guidelines for obtaining confidence bounds on a posterior density and suggest how to calibrate posterior probabilities in Bayesian testing.

Consider a prior density w on a d -dimensional real parameter θ indexing a collection of likelihoods p_θ . Suppose we have a random vector of data $X_n = (X_1, \dots, X_n)$ where the X_i 's are drawn independently from $p(\cdot | \theta) = p_{\theta_0}(\cdot)$. Denoting the outcomes by $x_n = (x_1, \dots, x_n)$, the posterior density is $w(\theta | x_n) = w(\theta) p_\theta(x_n) / m(x_n)$ where $m(\cdot)$ is the Bayesian marginal density for the data, that is, the mixture of distributions over θ , $m(x_n) = \int \mathbb{R}^d w(\theta) p_\theta(x_n) d\theta$.

Our main result is that when $p(\cdot | \theta)$ is an exponential family,

$$(1.1) \quad \int w(\theta_0 | x_n) p(x_n | \theta_0) dx_n = \frac{n^{d/2} |I(\theta_0)|^{1/2}}{2^{d/2} (2\pi)^{d/2}} \left\{ 1 + \frac{n}{L_1(\theta_0)} \right\} + o(n^{d/2-1}),$$

where θ_0 is the value of the parameter taken to be true, $I(\theta_0)$ is the Fisher information matrix and $L_1(\theta_0)$ is a quantity depending on the prior and its first two derivatives, but not on n . The quantity $L_1(\theta_0)$ is identified in the proof. Outside of very simple examples such as normal priors and normal likelihoods, it is very difficult to evaluate the mean, let alone the variance of a posterior, when the integration is over the sample space. However, results such as (1.1) become possible when the sample size n is allowed to increase.

The technique of proof for (1.1) can be adapted to give an asymptotic expression for the expected square of the posterior, namely:

$$(1.2) \quad \int [w(\theta_0 | x_n)]^2 dx_n | \theta_0 = \frac{n^d |I(\theta_0)|}{3^{d/2} (2\pi)^{d/2}} \left\{ 1 + \frac{n}{L_2(\theta_0)} \right\} + o(n^{d-1})$$

where $L_2(\theta_0)$ is another quantity depending only on the prior and its first two derivatives. Our methods generalize to give expansions with smaller error terms. Note that (1.1) and (1.2) constitute a frequentist assessment of the Bayesian's posterior. Indeed, if a Bayesian were to take an expected value of a posterior, for experimental design purposes for instance, the expectation would be with respect to the mixture density, $m(x_n) = \int w(\theta) p(x_n | \theta) d\theta$. This gives the trivial result $E^m w(\theta | X_n) = w(\theta)$.

The structure of this paper is as follows. In Section 2 we formally obtain (1.1) for exponential families. In Section 3 we formally obtain (1.2), also for exponential families. In Section 4 we discuss implications of our results for frequentist confidence bands around the posterior and for hypothesis testing. Proofs of the more technical results are gathered into the Appendices at the end.

2. Asymptotic bounds for the expected posterior

2.1 Notation

Our first task will be giving upper and lower bounds on $m(x_n)/d(x_n | \theta)$, which will be valid on sets whose probability tends to unity. As a consequence, we

will obtain a lower bound on the expected posterior and a corresponding upper bound that consists of two terms. The first is similar to the lower bound and the other is an error term which is typically negligible. To begin this program we introduce some formalities. We assume that the densities p_θ are twice continuously differentiable with respect to θ for almost every x and that there is a δ so that for each j and k from 1 to d we have that

$$(2.1) \quad E \sup_{\|\theta - \theta_0\| \leq \delta} \left| \frac{\partial^2}{\partial \theta_i \partial \theta_j} \log p(X_1 | \theta) \right|^2$$

is finite and for each i we have that

$$(2.2) \quad E \left| \frac{\partial}{\partial \theta_i} \log p(X_1 | \theta_0) \right|$$

is finite. Expectations here are with respect to the n -fold product of $p(\cdot | \theta_0)$, where θ_0 is a fixed value for θ . We write $p_{\theta_0}(\cdot) = p(\cdot | \theta_0)$ for convenience, and sometimes write E_{θ_0} for emphasis. We assume the family is soundly parameterised in the sense that convergence of a sequence of parameter values is equivalent to the weak convergence of the distributions they index. Denote the standardized score function by

$$(2.3) \quad \ell'_n(\theta) = \frac{1}{n} \nabla \log p_\theta(X^n),$$

and the empirical Fisher information by the $d \times d$ matrix

$$(2.4) \quad I^*(\theta) = \left(-\frac{1}{n} \sum_{k=1}^n \frac{\partial^2}{\partial \theta_i \partial \theta_j} \log p(X_k | \theta) \right)_{i,j=1,\dots,d}$$

We use a local argument on neighborhoods of the form

$$(2.5) \quad N(\theta_0, \alpha) = \{\theta | (\theta - \theta_0)^t I(\theta_0) (\theta - \theta_0) \leq \alpha^2\},$$

where $\alpha > 0$ and $I(\theta)$ is the true Fisher information at θ , assumed to be strictly positive and finite. For $\varepsilon, \delta > 0$ and satisfying $2\delta/(1 - \varepsilon) < \alpha$ we define 3 sets. The first controls the behavior of the posterior. It is

$$(2.6) \quad A_n(\alpha, \varepsilon, \theta_0) = \left\{ X^n | W(N(\theta_0, \alpha) | X^n) \geq \frac{1}{1 + \varepsilon} \right\},$$

where $W(\cdot | X^n)$ is the probability corresponding to the posterior density $w(\cdot | X^n)$. The second controls $I^*(\theta_0)$ on $N(\theta_0, \alpha)$. It is

$$(2.7) \quad \begin{aligned} B_n(\alpha, \varepsilon, \theta_0) = \{ X^n | & (1 - \varepsilon)(\theta - \theta_0)^t I(\theta_0) (\theta - \theta_0) \\ & \leq (\theta - \theta_0)^t I^*(\theta') (\theta - \theta_0) \\ & \leq (1 + \varepsilon)(\theta - \theta_0)^t I(\theta_0) (\theta - \theta_0) \\ & \text{for } \theta, \theta' \in N(\theta_0, \alpha) \}. \end{aligned}$$

The third controls the score:

$$(2.8) \quad C_n(\delta, \theta_0) = \{ X^n | \ell'_n(\theta_0)^t I^{-1}(\theta_0) \ell'_n(\theta_0) \leq \delta^2 \}.$$

2.2 Bounds on the density ratio

We use a version of Laplace's method to obtain bounds on $m(x^n)/p(x^n | \theta)$. Since we want to examine dependence of the expected posterior on the prior it is not enough to approximate $w(\theta)$ on $N(\theta_0, \alpha)$ by $w(\theta_0)$. Indeed, we use a second order Taylor expansion of the prior density. To control the error, we suppose there are functions $\nabla^2 \bar{w}(\theta_0, \alpha)$ and $\nabla^2 \underline{w}(\theta_0, \alpha)$ so that

$$(2.9) \quad \lim_{\alpha \rightarrow 0} \nabla^2 \bar{w}(\theta_0, \alpha) = \lim_{\alpha \rightarrow 0} \nabla^2 \underline{w}(\theta_0, \alpha) = \nabla^2 w(\theta_0)$$

and for $\theta, \theta' \in N(\theta_0, \alpha)$ we have the bound

$$(2.10) \quad \begin{aligned} (\theta - \theta_0)^t \nabla^2 \underline{w}(\theta_0, \alpha) (\theta - \theta_0) &\leq (\theta - \theta_0)^t \nabla^2 w(\theta') (\theta - \theta_0) \\ &\leq (\theta - \theta_0)^t \nabla^2 \bar{w}(\theta_0, \alpha) (\theta - \theta_0). \end{aligned}$$

Then, on $N(\theta_0, \alpha)$ we have upper and lower bounds on the prior density given by

$$(2.11a) \quad w(\theta) \leq w(\theta_0) + (\theta - \theta_0)^t \nabla w(\theta_0) + \frac{1}{2} (\theta - \theta_0)^t \nabla^2 \bar{w}(\theta_0, \alpha) (\theta - \theta_0),$$

and

$$(2.11b) \quad w(\theta) \geq w(\theta_0) + (\theta - \theta_0)^t \nabla w(\theta_0) + \frac{1}{2} (\theta - \theta_0)^t \nabla^2 \underline{w}(\theta_0, \alpha) (\theta - \theta_0).$$

We use (2.11a, b) to bound the density ratio by transforming to a quantity in $(\theta - u)$, where $u = \theta_0 + \frac{1}{1-\varepsilon} I^{-1}(\theta_0) \ell'_n(\theta_0)$. To state our bounds requires that we introduce some extra functions which occur in those bounds. For the sake of exposition we also define some related functions which will be used in the course of the proof. For the upper bound on the density ratio we use five functions:

$$(2.12a) \quad \begin{aligned} G_1 &= w(\theta_0) + \frac{1}{1-\varepsilon} \ell'_n(\theta_0)^t I^{-1}(\theta_0) \nabla w(\theta_0) \\ &\quad + \frac{1}{2(1-\varepsilon)^2} \ell'_n(\theta_0)^t I^{-1}(\theta_0) \nabla^2 \bar{w} I^{-1}(\theta_0) \ell'_n(\theta_0), \end{aligned}$$

$$(2.12b) \quad G_2 = (\theta - u)^t \left(\nabla w(\theta_0) + \frac{1}{1-\varepsilon} \nabla^2 \bar{w} I^{-1}(\theta_0) \ell'_n(\theta_0) \right),$$

$$(2.12c) \quad G_3 = \frac{1}{2} (\theta - u)^t \nabla^2 \bar{w}(\theta - u),$$

$$(2.12d) \quad G_4 = \nabla w(\theta_0) + \frac{1}{1-\varepsilon} \nabla^2 \bar{w} I^{-1}(\theta_0) \ell'_n(\theta_0),$$

$$(2.12e) \quad H_n = (1 + \varepsilon) e^{(n/2)(1-\varepsilon) \ell'_n(\theta_0)^t I^{-1}(\theta_0) \ell'_n(\theta_0)}.$$

Note that the sum $G_1 + G_2 + G_3$ is the right hand side of (2.11a). It will be seen that G_4 and H_n appear explicitly in the upper bound. For the lower bound we use five analogous functions:

$$(2.13a) \quad \tilde{G}_1 = w(\theta_0) + \frac{1}{1+\varepsilon} \ell'_n(\theta_0)^t I^{-1}(\theta_0) \nabla w(\theta_0)$$

$$+ \frac{1}{2(1+\varepsilon)^2} \ell'_n(\theta_0)^t I^{-1}(\theta_0) \nabla^2 \underline{w} I^{-1}(\theta_0) \ell'_n(\theta_0),$$

$$(2.13b) \quad \tilde{G}_2 = (\theta - u)^t \left(\nabla w(\theta_0) + \frac{1}{1+\varepsilon} \nabla^2 \underline{w} I^{-1}(\theta_0) \ell'_n(\theta_0) \right),$$

$$(2.13c) \quad \tilde{G}_3 = \frac{1}{2} (\theta - u)^t \nabla^2 \underline{w} (\theta - u),$$

$$(2.13d) \quad \tilde{G}_4 = \nabla w(\theta_0) + \frac{1}{1+\varepsilon} \nabla^2 \underline{w} I^{-1}(\theta_0) \ell'_n(\theta_0),$$

$$(2.13e) \quad \tilde{H}_n = (1-\varepsilon) e^{(n/2)(1+\varepsilon)} \ell'_n(\theta_0)^t I^{-1}(\theta_0) \ell'_n(\theta_0).$$

Analogously, the sum $\tilde{G}_1 + \tilde{G}_2 + \tilde{G}_3$ gives the right hand side of (2.11b).

Now we can state and prove our bounds on the density ratio.

PROPOSITION 2.1. (a) *Upper bound on $m(X^n)/p(X^n | \theta_0)$: Suppose $\delta/(1-\varepsilon) < \alpha/2$ and the prior density admits a second order Taylor expansion as in (2.11a). Then on the set $A_n \cap B_n \cap C_n$ we obtain the upper bound*

$$(2.14a) \quad \frac{m^n(X^n)}{p(X^n | \theta_0)} \leq \frac{H_n(2\pi)^{d/2}}{|n(1-\varepsilon)I(\theta_0)|^{1/2}} \cdot \left(G_1 + \frac{\text{tr} \nabla^2 \bar{w} I^{-1}(\theta_0)}{2n(1-\varepsilon)} + C_1(\theta_0) e^{-(n/16)\alpha^2(1-\varepsilon)} \right),$$

where

$$C_1(\theta_0) = 2^{d/2} \left(|G_1| + \left| \frac{2G_4^t I^{-1}(\theta_0) G_4}{n(1-\varepsilon)} \right|^{1/2} + \frac{\sqrt{3 \text{tr}[\nabla^2 \bar{w} I^{-1}(\theta_0) \nabla^2 \bar{w} I^{-1}(\theta_0)]}}{n(1-\varepsilon)} \right).$$

Here tr denotes the trace function of a matrix.

(b) *Lower bound on $m(X^n)/p(X^n | \theta_0)$: Suppose $\delta/(1+\varepsilon) < \alpha/2$ and the prior density admits a second order Taylor expansion as in (2.11b). Then on the set $B_n \cap C_n$ we have the lower bound*

$$(2.14b) \quad \frac{m_n(X^n)}{p(X^n | \theta_0)} \geq \frac{\tilde{H}_n(2\pi)^{d/2}}{|n(1+\varepsilon)I(\theta_0)|^{1/2}} \cdot \left(\tilde{G}_1 + \frac{\text{tr} \nabla^2 \underline{w} I^{-1}(\theta_0)}{2n(1+\varepsilon)} + C_2(\theta_0) e^{-(n/16)\alpha^2(1+\varepsilon)} \right),$$

where

$$C_2(\theta_0) = 2^{d/2} \left(|\tilde{G}_1| + \left| \frac{2\tilde{G}_4^t I^{-1}(\theta_0) \tilde{G}_4}{n(1+\varepsilon)} \right|^{1/2} + \frac{\sqrt{3 \text{tr}[\nabla^2 \underline{w} I^{-1}(\theta_0) \nabla^2 \underline{w} I^{-1}(\theta_0)]}}{n(1+\varepsilon)} \right).$$

Remark. In the proof of Theorem 2.1 in the Appendix A, it is shown that under the assumptions already stated at the beginning of this section the probabilities of the sets A_n , B_n and C_n are $o(1/n)$. Thus, from (2.14a, b) we will be able to obtain a characterization of $m_n(X^n)/p(X^n | \theta_0)$ which is exact enough to approximate the expected posterior asymptotically.

PROOF. See the Appendix A. \square

2.3 *Bounds on the posterior*

Next, we use the bounds in Proposition 2.1 to obtain bounds on the expected posterior, $E_{\theta_0} w(\theta_0 | X^n)$. First, we give a lower bound, then we give an upper bound of two parts. The first part of the upper bound is the same as the lower bound. The second part of the upper bound is an additional error term. The main factor in it is

$$J_n(\theta_0) = \int_{(B_n \cap C_n)^c} w(\theta_0 | x^n) p(x^n | \theta_0) dx^n.$$

For convenience we write $\beta = 2^{d/2} (2\pi)^{-d/2} |I(\theta_0)|^{1/2}$.

We are unable to demonstrate that J_n goes to zero in general. However, in Proposition 2.2 we give conditions under which J_n will go to zero when p_θ is a member of a parametric family in exponential form. We conjecture that it does in fact go to zero under much weaker conditions but have not been able to identify them.

To state Theorem 2.1, let

$$\begin{aligned} L_1(\theta_0) = & \frac{\nabla w(\theta_0)^t I^{-1}(\theta_0) \nabla w(\theta_0)}{2w^2(\theta_0)} - \frac{3 \operatorname{tr} \nabla^2 w(\theta_0) I^{-1}(\theta_0)}{4w(\theta_0)} \\ & - 2^{d/2} E[p_{2,\theta_0}(Z) e^{-Z^t Z/2}] \\ & - \frac{2^{d/2}}{w(\theta_0)} \nabla w^t(\theta_0) I^{-1/2}(\theta_0) E[Z p_{1,\theta_0}(Z) e^{-Z^t Z/2}], \end{aligned}$$

where Z is *Normal*(0, $I_{d \times d}$) and the polynomials $p_{1,\theta_0}(Z)$ and $p_{2,\theta_0}(Z)$ are given in terms of the cumulants of the distribution defined by the density p_{θ_0} . In the unidimensional parameter case they are the Hermite polynomials, see Feller (1971). The general definition for the multidimensional case is given in Bhattacharya and Rao (1986).

THEOREM 2.1. *Assume the conditions stated at the beginning of this section and in Proposition 2.1. Suppose X_i is continuous with positive definite Fisher information matrix $I(\theta)$ and has finite fourth moments. Also, assume that the characteristic function of X_1 is an element of L^p for some $p \geq 1$. Then we have the following bounds. Lower bound on the expected posterior:*

$$(2.15a) \quad \underline{\lim}_{n \rightarrow \infty} n \left\{ \frac{2^{d/2} (2\pi)^{d/2}}{n^{d/2} |I(\theta_0)|^{1/2}} \int w(\theta_0 | x^n) p(x^n | \theta_0) dx^n - 1 \right\} \geq L_1(\theta_0).$$

Upper bound on the expected posterior:

$$(2.15b) \quad \begin{aligned} \overline{\lim}_{n \rightarrow \infty} n \left\{ \frac{2^{d/2} (2\pi)^{d/2}}{n^{d/2} |I(\theta_0)|^{1/2}} \int w(\theta_0 | x^n) p(x^n | \theta_0) dx^n - 1 \right\} \\ \leq L_1(\theta_0) + \beta \overline{\lim}_{n \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \frac{J_n(\theta_0)}{n^{d/2-1}}, \end{aligned}$$

where the first limit on the right hand side of (2.15b) is over appropriately chosen sequences of $\alpha_n, \varepsilon_n, \delta_n \rightarrow 0$, satisfying $2\delta_n / (1 - \varepsilon_n) < \alpha_n$.

PROOF. See the Appendix A. \square

Our next result gives sufficient conditions for $J_n(\theta) \rightarrow 0$ for appropriately chosen sequences of $\alpha_n, \varepsilon_n, \delta_n \rightarrow 0$. Suppose the parametric family p_θ is of exponential family with d parameters in the natural parameterization. That is, write

$$(2.16) \quad p(x | \theta) = e^{\theta \cdot x - \psi(\theta)}.$$

We show that the extra error term in the upper bound (2.15b) is bounded above by e^{-c^n} for some positive c . Note that in an exponential family we can assume that the set B_n is the entire sample space since the second derivative of the log likelihood is not a function of the data x^n . Furthermore, the set C_n reduces to the set on which the maximum likelihood estimator is not close to the true value of the parameter. Thus our task is to bound quantities of the form

$$E_{\theta_0}[w(\theta_0 | X^n)1_{\{|\hat{\theta} - \theta_0| \geq \varepsilon\}}]$$

where $\hat{\theta}$ is the MLE of θ_0 .

PROPOSITION 2.2. *Suppose that p_θ is an exponential family with the natural parametrization. Then, if the natural parameter space has nonvoid interior, we have that there is a $\xi = \xi(\varepsilon) > 0$ so that*

$$(2.17) \quad E_{\theta_0}[w(\theta_0 | X^n)1_{\{|\hat{\theta} - \theta_0| > \varepsilon\}}] \leq e^{-\xi^n}.$$

PROOF. We have that

$$p(x^n | \theta) = \exp \left\{ \theta \cdot \sum_{i=1}^n x_i + n\psi(\theta) \right\},$$

where $\theta = (\theta_1, \dots, \theta_d)$ and $x_i = (x_{i1}, \dots, x_{id})$ for $i = 1, \dots, n$. Consequently, for fixed n we obtain

$$(2.18) \quad \int w(\theta) \frac{p(x^n | \theta)}{p(x^n | \theta_0)} d\theta \geq \int_{\tilde{E}} w(\theta) \exp \left\{ (\theta - \theta_0) \sum_{i=1}^n x_i + n[\psi(\theta) - \psi(\theta_0)] \right\} d\theta,$$

where $\tilde{E} \equiv \tilde{E}(\varepsilon', \theta_0, \sum_{i=1}^n x_i) \subset \Omega$ is defined as follows. For $\varepsilon' > 0$, we require $|\theta_i - \theta_{oi}| < \varepsilon'$ and for given $\sum_{i=1}^n x_i$, we require $(\theta_i - \theta_{oi}) \sum_{j=1}^n x_{ij} \geq 0$, where $\theta_0 = (\theta_{o1}, \dots, \theta_{od})$. Since $\tilde{E}(\varepsilon', \theta_0, \sum_{i=1}^n x_i)$ depends on $\sum_{i=1}^n x_i$ only on the signs of its d entries, there are 2^d possible sets. Denote them by $E_1(\varepsilon', \theta_0), \dots, E_{2^d}(\varepsilon', \theta_0)$. Now we can bound the right hand side of (2.18) from below by

$$(2.19) \quad \inf_{|\theta - \theta_0| < \varepsilon'} w(\theta) \min_{1 \leq i \leq 2^d} \int_{E_i(\varepsilon', \theta_0)} \exp\{n[\psi(\theta) - \psi(\theta_0)]\} d\theta.$$

Since w is continuous and ψ is convex, we know that for given $\eta > 0$, and $\delta > 0$, there is $\varepsilon' > 0$ so small that

$$(2.20) \quad \psi(\theta) - \psi(\theta_0) \geq -\eta \quad \text{and} \quad w(\theta) - w(\theta_0) \geq -\delta,$$

if $\max_{1 \leq i \leq d} |\theta_i - \theta_{0i}| < \varepsilon'$. Thus (2.19) and the left hand side of (2.18) is bounded below by

$$(2.21) \quad [w(\theta_0) - \delta](\varepsilon')^d e^{-n\eta}.$$

We use (2.21) to obtain the proposition. The right hand side of (2.17) is

$$(2.22) \quad w(\theta_0) \int_{|\hat{\theta} - \theta_0| > \varepsilon} \left\{ \int w(\theta) \frac{p(x^n | \theta)}{p(x^n | \theta_0)} d\theta \right\}^{-1} p(x^n | \theta_0) dx^n \\ \leq \frac{w(\theta_0) e^{n\eta} P_{\theta_0}(|\hat{\theta} - \theta_0| \geq \varepsilon)}{[w(\theta_0) - \delta](\varepsilon')^d}.$$

Since the natural parameter space is convex and has nonvoid interior, p_θ has a moment generating function which is finite on an neighborhood of zero. So, for given $\varepsilon > 0$, there is a $c(\varepsilon) > 0$, so that $P_{\theta_0}(|\hat{\theta} - \theta_0| \geq \varepsilon) \leq \exp\{-nc(\varepsilon)\}$. Now for given $\varepsilon > 0$, it is enough to choose $\varepsilon' > 0$ and δ so small that (2.49) is satisfied for some $\eta > 0$ and $\delta > 0$ which also satisfies $c(\varepsilon) - \eta > 0$. \square

3. Asymptotic bounds for the expected square of the posterior

In this section we use adapt the proof of Theorem 2.1 to establish an asymptotic expansion for the expected square of the posterior. First, we define analogs of the quantities appearing in Theorem 2.1 and Proposition 2.2. Let

$$(3.1a) \quad L_2(\theta_0) = \frac{\nabla w(\theta_0)^t I^{-1}(\theta_0) \nabla w(\theta_0)}{w^2(\theta_0)} - \frac{4 \operatorname{tr} \nabla^2 w(\theta_0) I^{-1}(\theta_0)}{3w(\theta_0)} \\ - 3^{d/2} E[p_{2,\theta_0}(Z) e^{-Z^t Z}] \\ - \frac{2(3^{d/2})}{w(\theta_0)} \nabla w^t(\theta_0) I^{-1/2}(\theta_0) E[Z p_{1,\theta_0}(Z) e^{-Z^t Z}],$$

where Z is $Normal(0, I_{d \times d})$ distributed. Also, write

$$(3.1b) \quad \tilde{\beta} = \frac{|I(\theta_0)|}{3^{d/2}(2\pi)^d} \quad \text{and} \quad \tilde{J}_n(\theta_0) = \int_{(B_n \cap C_n)^c} [w(\theta_0 | x^n)]^2 p(x^n | \theta_0) dx^n.$$

Now we have the following theorem, and, as before, we control the error term $\tilde{\beta} \tilde{J}_n(\theta_0)$ in a separate result.

THEOREM 3.1. *Under the assumptions of Theorem 2.1, we have the following bounds:*

$$(3.2a) \quad \lim_{n \rightarrow \infty} n \left\{ \frac{3^{d/2}(2\pi)^d}{n^d |I(\theta_0)|} \int [w(\theta_0 | x^n)]^2 p(x^n | \theta_0) dx^n - 1 \right\} \geq L_2(\theta_0)$$

and

$$(3.2b) \quad \overline{\lim}_{n \rightarrow \infty} n \left\{ \frac{3^{d/2}(2\pi)^d}{n^d |I(\theta_0)|} \int [w(\theta_0 | x^n)]^2 p(x^n | \theta_0) dx^n - 1 \right\} \\ \leq L_2(\theta_0) + \tilde{\beta} \overline{\lim}_{n \rightarrow \infty} \overline{\lim} \frac{\tilde{J}_n(\theta_0)}{n^{d-1}},$$

where the first limit on the right hand side of (3.2b) is over appropriately chosen sequences of $\alpha_n, \varepsilon_n, \delta_n \rightarrow 0$, satisfying $2\delta_n/(1 - \varepsilon_n) < \alpha_n$.

PROOF. The proof is a straightforward modification of the proof of Theorem 2.1; it is given in the Appendix B. \square

The next result gives conditions under which $\tilde{J}_n(\theta) \rightarrow 0$ when p_θ is of exponential family as in (2.16). As before, the sets B_n and C_n reduce to the set on which the maximum likelihood estimator is far from the true value of the parameter so our task is to bound $E_{\theta_0} w(\theta_0 | X^n)^2 \chi_{\{|\hat{\theta} - \theta_0| \geq \varepsilon\}}$, where $\hat{\theta}$ is the MLE of θ_0 .

PROPOSITION 3.1. *Under the conditions of Proposition 2.2, we have that there is a $\xi = \xi(\varepsilon) > 0$ so that*

$$(3.3) \quad E_{\theta_0} [w(\theta_0 | X^n)^2 \chi_{\{|\hat{\theta} - \theta_0| > \varepsilon\}}] \leq e^{-\xi n}.$$

PROOF. This is a straightforward modification of the proof of Proposition 2.2. \square

4. Implications of the results

The original motivation for Theorem 2.1 arose from the search for noninformative priors. Bernardo (1979) proposed the use of what he called reference priors, obtained by maximizing the relative entropy distance between a prior and its corresponding posterior over a large class of possible priors. Clarke and Sun (1997) performed an analogous optimization using the Chi-squared distance in place of the relative entropy. In this latter context, the $O(1/n)$ terms in the expansion of Theorem 2.1 generate a functional which can be optimized so as to yield Chi-squared reference priors proportional to the inverse of the Fisher information. Since this application has been pursued at length elsewhere, we turn instead to implications of Theorems 2.1 and 3.1 more generally.

4.1 *Heuristic derivation of confidence bands for the expected posterior*

In Sections 2 and 3 formal asymptotic expansions for $E_{\theta_0} w(\theta_0 | X^n)$ and $E_{\theta_0} [w(\theta_0 | X^n)]^2$ were given for exponential families. These results lead to conjectures about the form of the expansions when the argument of the posterior density is not the true value of the parameter. Indeed, this case can be reduced to the case where the argument is θ_0 . Consider the identities

$$(4.1a) \quad E_{\theta_0} [w(\theta | X^n)] \\ = \frac{w(\theta)}{w(\theta_0)} E_{\theta_0} \left[w(\theta_0 | X^n) \exp \left(-n \left\{ \frac{1}{n} \log \left[\frac{p(x^n | \theta_0)}{p(x^n | \theta)} \right] \right\} \right) \right],$$

and

$$(4.1b) \quad E_{\theta_0} [w(\theta | X^n)]^2 \\ = \left\{ \frac{w(\theta)}{w(\theta_0)} \right\}^2 E_{\theta_0} \left[w(\theta_0 | X^n)^2 \exp \left(-2n \left\{ \frac{1}{n} \log \left[\frac{p(x^n | \theta_0)}{p(x^n | \theta)} \right] \right\} \right) \right].$$

By the law of large numbers, the exponent $n^{-1} \log[p(x^n | \theta_0)/p(x^n | \theta)]$ converges in various modes to the relative entropy between p_{θ_0} and p_{θ} , which we write as

$$D(\theta_0 \| \theta) = \int p(x^n | \theta_0) \log \frac{p(x^n | \theta_0)}{p(x^n | \theta)} dx^n.$$

Making this substitution heuristically in (4.1a,b) gives

$$(4.2a) \quad E_{\theta_0} w(\theta | X^n) \approx \left[\frac{w(\theta)}{w(\theta_0)} \right] e^{-nD(\theta_0 \| \theta)} E_{\theta_0} [w(\theta_0 | X^n)],$$

and

$$(4.2b) \quad E_{\theta_0} w(\theta | X^n)^2 \approx \left[\frac{w(\theta)}{w(\theta_0)} \right]^2 e^{-2nD(\theta_0 \| \theta)} E_{\theta_0} [w(\theta_0 | X^n)]^2$$

for large n . Now, using Theorem 2.1 and Theorem 3.1 we have

$$(4.3a) \quad E_{\theta_0} w(\theta | X^n) \approx \left[\frac{w(\theta)}{w(\theta_0)} \right] e^{-nD(\theta_0 \| \theta)} \sqrt{\frac{nI(\theta_0)}{4\pi}}$$

and

$$(4.3b) \quad E_{\theta_0} w(\theta | X^n)^2 \approx \left[\frac{w(\theta)}{w(\theta_0)} \right]^2 e^{-2nD(\theta_0 \| \theta)} \frac{nI(\theta_0)}{\sqrt{32\pi}},$$

leaving out the terms of order $O(1/n)$ and smaller. For the unidimensional case, $d = 1$ and we note that by expressing the variance as the mean square minus the square mean we have

$$(4.4) \quad \text{Var}_{\theta_0}(w(\theta | X^n)) \approx \left[\frac{w(\theta)}{w(\theta_0)} \right]^2 e^{-2nD(\theta_0 \| \theta)} \left(\frac{1}{\sqrt{3}} - \frac{1}{2} \right) \frac{n|I(\theta_0)|}{2\pi}.$$

If the details justifying (4.2a, b) can be supplied then one can obtain a confidence band for the expected posterior $E_{\theta_0} w(\theta | X^n)$. Note that the expected value

considers all the data that might have been got whereas the posterior, $w(\theta | X^n)$, only uses the actual set of data obtained.

It is interesting to note that the left hand side of (4.3a) is a probability density for θ : It is positive and integrates to unity. The right hand side is nearly a density also. The discrepancy arises from the neglect of terms that are $O(1/n)$ and smaller. Indeed, $D(\theta_0 || \theta)$ can be Taylor expanded at θ_0 to see that for θ near θ_0

$$D(\theta_0 || \theta) \approx \frac{1}{2}(\theta - \theta_0)^2 I(\theta_0)$$

because $D(\theta_0 || \theta_0) = D'(\theta_0 || \theta_0) = 0$. By Laplace's method, see De Bruijn (1961), it is straightforward to verify that

$$(4.5) \quad \int w(\theta) e^{-nD(\theta_0 || \theta)} d\theta \approx w(\theta_0) \sqrt{\frac{2\pi}{nI(\theta_0)}}$$

as $n \rightarrow \infty$. Comparison of (4.3a) and (4.5) suggests defining

$$(4.6) \quad g_n(\theta_0, \theta) = \frac{w(\theta)}{w(\theta_0)} \sqrt{\frac{nI(\theta_0)}{2\pi}} e^{-nD(\theta_0 || \theta)}.$$

Now, (4.3a, b) become

$$E_{\theta_0}[w(\theta | X^n)] \approx \frac{1}{\sqrt{2}} g_n(\theta_0, \theta)$$

and

$$E_{\theta_0}[w(\theta | X^n)^2] \approx \frac{1}{\sqrt{3}} g_n(\theta_0, \theta)^2.$$

That is, the near-density g_n encapsulates in an asymptotic sense the expectation of the posterior, apart from constant factors independent of the prior and parametric family. These factors $1/\sqrt{2}$ and $1/\sqrt{3}$ may arise from the fact that most location estimators, such as the maximum likelihood estimator are asymptotically Chi-square, a fact which was neglected in the application of Laplace's method since the expansion was around θ_0 rather than around an estimator for θ .

Now consider trying to estimate $E_{\theta_0}[w(\theta | X^n)]$, the expected posterior when θ_0 is true, by $w(\theta | X^n)$. Markov's inequality gives

$$P_{\theta_0}(|w(\theta | X^n) - E_{\theta_0}[w(\theta | X^n)]| \geq \epsilon) \leq \frac{\text{Var}_{\theta_0}[w(\theta | X^n)]}{\epsilon^2}.$$

So, it is seen from (4.4) that

$$(4.7) \quad P_{\theta_0}(|w(\theta | X^n) - E_{\theta_0}[w(\theta | X^n)]| \leq \epsilon) \geq 1 - \alpha$$

can be achieved for fixed confidence level $1 - \alpha$ by choosing

$$[\epsilon_n(\theta_0, \theta)]^2 = (1/\sqrt{3} - 1/2)[g_n(\theta_0, \theta)]^2.$$

That is, with this choice of $\epsilon_n(\theta_0, \theta)$,

$$(4.8) \quad w(\theta | x^n) \pm \frac{1}{\sqrt{1-\alpha}} \left(\frac{1}{\sqrt{3}} - \frac{1}{2} \right)^{1/2} g_n(\theta_0, \theta)$$

provides an approximate, asymptotic $1-\alpha$ confidence band for $E_{\theta_0}[w(\theta | X^n)]$ if one replaces θ_0 in (4.8) by an estimator such as the maximum likelihood estimator. Such a confidence band indicates the reliability of inferences from the posterior density by giving a range of nearby densities, based on data near x^n that might have been obtained. (In principle, one might get a better confidence band if it were established that the posterior density, for each θ was asymptotically normal with mean $E_{\theta_0}[w(\theta | X^n)]$ and variance $\text{Var}_{\theta_0}[w(\theta | X^n)]$ times some function of n , in the usual frequentist sense.)

4.2 *A frequentist refinement of Bayesian testing*

Consider testing $H_0 : \theta \in \Omega_0$ versus $H_1 : \theta \in \Omega_1$, where both hypotheses are the closures of open bounded sets in the parameter space. A Bayes test is performed by calculating $W(\Omega_0 | X^n)$, or, equivalently, the posterior odds ratio provided $\Omega_0 = \Omega_1^c$ where the complement is taken within the parameter space. It is well known that this procedure is Bayes optimal under generalized zero-one loss, see Casella and Berger (1990). A frequentist test, by contrast, specifies a rejection region say A , with the property that $\sup_{\theta \in \Omega_0} P_\theta(A) \leq \alpha$, where α is an upper bound on the probability of type I error. For frequentist optimality, one ensures $P_\theta(A)$ is as large as possible for $\theta \in \Omega_1$.

Kass and Raftery (1995) credit Good with the proposal that Bayes and frequentist testing can be partially reconciled by using the posterior probability as a test statistic. To see how this might work, note if the posterior probability is used to define a rejection region then one would accept H_0 when $W(\Omega_0 | x^n) \geq \beta$ where β is a threshold value. A frequentist would bound the power of this test by noting that, for $\theta \in \Omega_1$,

$$(4.9) \quad P_\theta(W(\Omega_0 | X^n) \geq \beta) \leq \frac{1}{\beta^2} E_\theta \left\{ \int_{\Omega_0} w(\theta' | X^n) d\theta' \right\}^2,$$

by Markov's inequality. Multiplying and dividing $w(\theta' | X^n)$ by $\sqrt{w(\theta')}$ and applying the Cauchy-Schwartz inequality gives that the expectation in (4.9) is bounded by

$$(4.10) \quad \int_{\Omega_0} w(\theta') d\theta' E_\theta \left\{ \int_{\Omega_0} \frac{w(\theta' | X^n)^2}{w(\theta')} d\theta' \right\}.$$

Noting that the first factor in (4.10) is bounded by unity, Fubini's theorem permits the interchange of expectation and integration so that (4.3b) gives

$$(4.11) \quad \frac{KnI(\theta)}{[\beta w(\theta)]^2} \int_{\Omega_0} w(\theta') e^{-2nD(\theta \parallel \theta')} d\theta',$$

as an approximate upper bound for (4.9), where K is a positive constant that can be determined. It is seen that as $n \rightarrow \infty$, the right hand side of (4.11) is of order $O(e^{-n\gamma})$ for some $\gamma > 0$. Apart from the value of γ this is the same rate as is achieved by the Neyman-Pearson test in the simple versus simple case, so it cannot be improved substantially.

Despite its Bayes optimality, a Bayes test is incomplete in the sense that rejecting H_0 when $W(\Omega_0 | X^n)$ is too small does not take into account data sets near X^n that might have given a different value for the posterior probability. In short, the Bayesian neglects the sampling distribution of $W(\Omega_0 | X^n)$. To remedy this, we suggest that H_0 should only be accepted when the posterior exceeds a threshold determined by use of the sampling distribution, to wit, accept H_0 when

$$(4.12) \quad W(\Omega_0 | X^n) \geq \sup_{\theta \in \Omega_0} \{E_\theta W(\Omega_0 | X^n) + 3\sqrt{\text{Var}_\theta[W(\Omega_0 | X^n)]}\}$$

and reject H_0 when

$$(4.13) \quad W(\Omega_0 | X^n) \leq \inf_{\theta \in \Omega_0} \{E_\theta W(\Omega_0 | X^n) - 3\sqrt{\text{Var}_\theta[W(\Omega_0 | X^n)]}\}.$$

Expression (4.3a) gives a heuristic asymptotic expansion for the expected posterior probability. We have been unable to derive an analogous expansion for the variance of the posterior probability. In between (4.12) and (4.13) the data cannot be regarded as sufficiently conclusive as to justify either H_0 or H_1 .

In effect, this procedure uses a frequentist criterion, power, with frequentist near optimality in the sense of the power function to set a threshold for a Bayes test. As a consequence, this test is simultaneously Bayes optimal and frequentist ‘good’. The decision to accept or reject the null is based on a quantity equivalent to the posterior odds ratio which minimizes the posterior risk and has a well defined rejection region with power function bounded below by a function of the form $1 - O(e^{-\gamma n})$, as $n \rightarrow \infty$.

Finally, we examine the case that H_0 is a dimensional reduction from H_1 . That is, for example, $H_0 : \mu_1, \dots, \mu_{p-1} \in \mathbb{R}, \mu_p = \mu_0$, and $H_1 : \mu_1, \dots, \mu_p \in \mathbb{R}$. Here μ_0 is a fixed constant. The usual Bayes factor would be

$$(4.14) \quad W(\Omega_0 | X^n) / W(\Omega_0^c | X^n).$$

It is difficult even to conjecture an asymptotic form for the mean and variance of (4.14). We can nevertheless partially identify the behavior of the numerator or denominator separately. This does not use the present results, but does help to understand how one might use the posterior probability of a hypothesis as a test statistic.

Parallel to (4.12) and (4.13), we would like to find forms for $E_\mu W(\Omega_0 | X^n)$ and $\text{Var}_\mu[W(\Omega_0 | X^n)]$ where $\mu = (\mu_1, \dots, \mu_p) \in \mathbb{R}^p$ assumed to be true. The denominator may be dependent statistically, but dealing with the numerator is enough for testing purposes.

Write the overall prior on μ as

$$w(\mu) = w(\tilde{\mu}^{p-1})[\alpha w_d(\mu_p) + (1 - \alpha)w_c(\mu_p)],$$

where $0 < \alpha < 1$ is a fixed constant, $\tilde{\mu}^{p-1} = (\mu_1, \dots, \mu_{p-1})$, w_c is a continuous prior for μ_p with support \mathbb{R} , and w_d assigns mass 1 to $\mu_p = \mu_0$. Write $m_d(X^n)$ and $m_c(X^n)$ to mean the mixture of the likelihoods $p(X^n | \mu)$ with respect to $w_d(\mu) = w(\tilde{\mu}^{p-1})w_d(\mu_p)$ and $w_c(\mu) = w(\tilde{\mu}^{p-1})w_c(\mu_p)$. Now, the overall mixture is $m(X^n) = \alpha m_d(X^n) + (1 - \alpha)m_c(X^n)$ and we have

$$(4.15) \quad w(\mu | X^n) = \lambda(X^n)w_d(\mu | X^n) + [1 - \lambda(X^n)]w_c(\mu | X^n),$$

where $w_d(\mu | X^n)$ and $w_c(\mu | X^n)$ are the posteriors formed from $w_d(\mu)$ and $w_c(\mu)$, respectively, and $\lambda(X^n) = \alpha m_d(X^n)/m(X^n)$.

Since $W_d(\Omega | X^n) = 1$ and $W_c(\Omega_0 | X^n) = 0$, integrating in (4.15) gives

$$(4.16) \quad W(\Omega_0 | X^n) = \lambda(X^n) = \left\{ 1 + \left(\frac{1 - \alpha}{\alpha} \right) \frac{m_c(X^n)}{m_d(X^n)} \right\}^{-1}.$$

The main quantity in (4.16) is the density ratio in the denominator. Suppose we fix an element of Ω_0 defined by $\tilde{\mu}^{p-1}$ and $\mu_p = \mu_0$. Then $m_c(X^n)/m_d(X^n)$ is

$$(4.17) \quad \frac{\int w_c(\mu)p(X^n | \mu)d\mu}{\int w(\tilde{\mu}^{p-1})p(X^n | \tilde{\mu}^{p-1}, \mu_0)d\tilde{\mu}^{p-1}} \frac{p(X^n | \tilde{\mu}^{p-1}, \mu_0)}{p(X^n | \tilde{\mu}^{p-1}, \mu_0)}$$

$$= \frac{m_c(X^n)}{p(X^n | \tilde{\mu}^{p-1}, \mu_0)} \frac{p(X^n | \tilde{\mu}^{p-1}, \mu_0)}{\int w(\tilde{\mu}^{p-1})p(X^n | \tilde{\mu}^{p-1}, \mu_0)d\tilde{\mu}^{p-1}}.$$

The two density ratios in (4.17) have behavior partially given by Theorem 2.1, Clarke and Barron (1990). This result gives conditions under which

$$(4.18) \quad \log \left\{ \frac{p(X^n | \mu)}{m(X^n)} \right\} = \frac{d}{2} \log \left(\frac{n}{2\pi} \right) + \frac{1}{2} S J^{-1}(\mu) S - \log[w(\mu)]$$

$$+ \frac{1}{2} \log |J(\mu)| + o_p(1),$$

where d is the dimension of μ , $S = \frac{1}{\sqrt{n}} \nabla \log p(X^n | \mu)$, and J is the Fisher information matrix. Using (4.18) for $d = p$ and $d = p - 1$ gives approximations for the two densities in (4.17). Substituting those approximations into (4.17) and then (4.16) gives a term of order $O_p(n^{-1/2})$. Thus, the expectation of (4.16) can be approximated as

$$(4.19) \quad E_\mu[W(\Omega_0 | X^n)] \approx \left\{ 1 + \left(\frac{1 - \alpha}{\alpha} \right) 0_p(n^{-1/2}) \right\}^{-1} \rightarrow 1.$$

If we consider $\mu \in \Omega_0$ then (4.16) still holds and if $\mu = (\mu_1, \dots, \mu_{p-1}, \mu'_p)$, $\mu'_p \neq \mu_0$, we get

$$(4.20) \quad \frac{m_c(X^n)}{p(X^n | \tilde{\mu}^{p-1}, \mu'_p)} \frac{p(X^n | \tilde{\mu}^{p-1}, \mu'_p)}{p(X^n | \tilde{\mu}^{p-1}, \mu_0)} \frac{p(X^n | \tilde{\mu}^{p-1}, \mu_0)}{\int w(\tilde{\mu}^{p-1})p(X^n | \tilde{\mu}^{p-1}, \mu_0)d\tilde{\mu}^{p-1}}$$

in place of (4.17). Again, using (4.18) in the first and third factors of (4.20) (cases $d = p, p - 1$ as before) and using

$$(4.21) \quad \frac{p(X^n | \tilde{\mu}^{p-1}, \mu'_p)}{p(X^n | \tilde{\mu}^{p-1}, \mu_0)} = \exp[n\hat{D}(\tilde{\mu}^{p-1}, \mu'_p || \tilde{\mu}^{p-1}, \mu_0)],$$

where \hat{D} is the empirical relative entropy between distributions with the indicated parameter values, the middle factor of (4.20) gives an asymptotic approximation for use in (4.16). This gives the expression

$$(4.22) \quad E_{(\tilde{\mu}^{p-1}, \mu'_p)}[W(\Omega_0 | X^n)] \approx \left\{ 1 + \left(\frac{1 - \alpha}{\alpha} \right) 0_p(e^{n\hat{D}} n^{-1/2}) \right\}^{-1} \rightarrow 0.$$

for any $(\tilde{\mu}^{p-1}, \mu'_p)$.

In closing, we note that a more careful analog would give more refined asymptotic forms for (4.19) and (4.22), and might give similarly refined forms for the variances of $W(\Omega_0 | X^n)$. Note that the analysis here is in probability only, even though (4.18) holds in an L^1 sense also. Extending the present heuristics would justify the decision rule suggested by (4.12) and (4.13).

Acknowledgements

The authors thank an anonymous referee for suggestions that improved an earlier version of this manuscript.

Appendix A: Proofs from Section 2

In this Appendix we have recorded proofs the two main technical results in Section 2, namely Proposition 2.1 and Theorem 2.1.

PROOF OF PROPOSITION 2.1. First we obtain (2.14a). Note that as in Clarke and Barron (1990) p. 463 we have on $A_n \cap B_n$ that

$$(A.1) \quad \frac{m_n(X^n)}{p_{\theta_0}(X^n)} \leq (1 + \varepsilon) e^{(n/2)(1-\varepsilon)\ell'_n(\theta_0)^t I^{-1}(\theta_0)\ell'_n(\theta_0)} \int_{N(\theta_0, \alpha)} e^{-(n/2)(1-\varepsilon)(\theta-u)^t I(\theta_0)(\theta-u)} w(\theta) d\theta,$$

where $u = \theta_0 + \frac{1}{1-\varepsilon} I^{-1}(\theta_0)\ell'_n(\theta_0)$. Now we have the left hand side of (A.1) is bounded by

$$(A.2) \quad \frac{m_n(X^n)}{p(X^n | \theta_0)} \leq H_n \left(\sum_{i=1}^3 \int_{\mathbb{R}^d} e^{-(n(1-\varepsilon)/2)(\theta-u)^t I(\theta_0)(\theta-u)} G_i d\theta - \sum_{i=1}^3 \int_{N(\theta_0, \alpha)^c} e^{-(n(1-\varepsilon)/2)(\theta-u)^t I(\theta_0)(\theta-u)} G_i d\theta \right).$$

The summands in the first summation of the right hand side of (A.2) can be evaluated. The first gives

$$(A.3) \quad \frac{H_n G_1 (2\pi)^{d/2}}{|n(1-\varepsilon)I(\theta_0)|^{1/2}},$$

the second is zero, and the third is

$$(A.4) \quad \frac{H_n}{2} \frac{(2\pi)^{d/2}}{|n(1-\varepsilon)I(\theta_0)|^{1/2}} E[Z^t \nabla^2 \bar{w} Z],$$

where $Z \sim \text{Normal}(0, \frac{1}{n(1-\varepsilon)} I^{-1}(\theta_0))$. We note that the expectation in (A.4) is

$$\begin{aligned} E[\text{tr}(Z^t \nabla^2 \bar{w} Z)] &= E[\text{tr}(\nabla^2 \bar{w} Z Z^t)] = \text{tr}(\nabla^2 \bar{w} E Z Z^t) \\ &= \text{tr} \left[\nabla^2 \bar{w} \frac{I^{-1}(\theta_0)}{n(1-\varepsilon)} \right] = \frac{1}{n(1-\varepsilon)} \text{tr}[\nabla^2 \bar{w} I^{-1}(\theta_0)]. \end{aligned}$$

So, the third term in the first summation on the right hand side of (A.2) is

$$(A.5) \quad \frac{H_n}{2} \frac{(2\pi)^{d/2}}{|n(1-\varepsilon)I(\theta_0)|^{1/2}} \cdot \frac{1}{n(1-\varepsilon)} \text{tr}(\nabla^2 \bar{w} I^{-1}(\theta_0)).$$

Next, we control the terms in the second summation on the right hand side of (A.2). The norm is defined by $I(\theta_0)$; θ is in $N(\theta_0, \alpha)^c$; and we have that $\delta/(1-\varepsilon) < \alpha/2$. Consequently, we have

$$(A.6) \quad \|\theta - u\| \geq \|\theta - \theta_0\| - \frac{1}{1-\varepsilon} \|I^{-1}(\theta_0) \ell'_n(\theta_0)\| \geq \frac{\alpha}{2}$$

on C_n .

The first term in the second sum in (A.2) is controlled as follows. By (A.6), we have that

$$(A.7) \quad e^{-(n/2)(1-\varepsilon)(\theta-u)^t I(\theta_0)(\theta-u)} \leq e^{-(n/4)(1-\varepsilon)(\theta-u)^t I(\theta_0)(\theta-u)} e^{-(n/16)\alpha^2(1-\varepsilon)}.$$

Now, we can bound the first term in the second sum by using this last inequality and enlarging the domain of integration. The result is

$$(A.8) \quad H_n |G_1| \frac{(2\pi)^{d/2} 2^{d/2}}{|n(1-\varepsilon)I(\theta_0)|^{1/2}} e^{-(n/16)\alpha^2(1-\varepsilon)}.$$

By enlarging the domain of integration and applying the Cauchy-Schwartz inequality, the second term in the second sum in (A.2) is bounded above by

$$(A.9) \quad \begin{aligned} H_n e^{-(n/16)(1-\varepsilon)\alpha^2} \int_{N_\varepsilon} \left| \left(\nabla w(\theta) + \frac{\nabla^2 \bar{w} I^{-1}(\theta_0) \ell'_n(\theta_0)}{1-\varepsilon} \right)^t (\theta - u) \right| \\ \cdot e^{-(n/4)(1-\varepsilon)(\theta-u)^t I(\theta_0)(\theta-u)} d\theta \\ \leq H_n e^{-(n/16)(1-\varepsilon)\alpha^2} \frac{(2\pi)^{d/2} 2^{d/2}}{|n(1-\varepsilon)I(\theta_0)|^{1/2}} \sqrt{E Z^t G_4 G_4^t Z}, \end{aligned}$$

where $Z \sim Normal(0, \frac{2}{n(1-\varepsilon)}I^{-1}(\theta_0))$. The expectation in (A.9) can be evaluated: Rearranging under trace gives

$$(A.10) \quad EZ^t G_4 G_4^t Z = \frac{2}{n(1-\varepsilon)} \text{tr}[G_4 G_4^t I^{-1}(\theta_0)] = \frac{2}{n(1-\varepsilon)} G_4^t I^{-1}(\theta_0) G_4.$$

Using (A.10) in (A.9) gives the bound

$$(A.11) \quad H_n e^{-(n/16)(1-\varepsilon)\alpha^2} \frac{(2\pi)^{d/2} 2^{d/2}}{|n(1-\varepsilon)I(\theta_0)|^{1/2}} \sqrt{\frac{2}{n(1-\varepsilon)} G_4^t I^{-1}(\theta_0) G_4}.$$

To finish obtaining the upper bound note that the third summand in the second summation in (A.2) is bounded above by

$$(A.12) \quad e^{-(n/16)(1-\varepsilon)\alpha^2} \frac{H_n}{2} \frac{(2\pi)^{d/2} 2^{d/2}}{|n(1-\varepsilon)I(\theta_0)|^{1/2}} \sqrt{E(Z^t \nabla^2 \bar{w} Z)^2},$$

where $Z \sim Normal(0, \frac{2}{n(1-\varepsilon)}I^{-1}(\theta_0))$. To evaluate the expectation in (A.12) note that for any $d \times d$ matrix T ,

$$E(Z^t T Z) = \text{tr} \left[\frac{2}{n(1-\varepsilon)} T I^{-1}(\theta_0) \right],$$

and

$$\text{Var}(Z^t T Z) = 2 \text{tr} \left[\frac{4}{n^2(1-\varepsilon)^2} T I^{-1}(\theta_0) T I^{-1}(\theta_0) \right].$$

So, (A.12) is bounded from above by

$$(A.13) \quad H_n \frac{(2\pi)^{d/2} 2^{d/2} \sqrt{3}}{|n(1-\varepsilon)I(\theta_0)|^{1/2} n(1-\varepsilon)} \cdot e^{-(n/16)(1-\varepsilon)\alpha^2} \sqrt{\text{tr} \nabla^2 \bar{w} I^{-1}(\theta_0) \nabla^2 \bar{w} I^{-1}(\theta_0)}.$$

Since all terms in the second sum in (A.2) (namely, (A.8), (A.11) and (A.13)) are of lower order than the terms in the first sum in (A.2) (namely, (A.3), zero and (A.5)) we have that the upper bound (2.14a) holds with $C_1(\theta_0)$ as defined.

Next, we derive (2.14b). From Clarke and Barron (1990) p. 463, we observe that

$$(A.14) \quad \frac{m_n(X^n)}{p(X^n | \theta_0)} \geq e^{(n/2(1+\varepsilon))\ell'_n(\theta_0)^t I(\theta_0)^{-1} \ell'_n(\theta_0)} \int_{N(\theta_0, \alpha)} e^{-((1+\varepsilon)n/2)(\theta-u)^t I(\theta_0)(\theta-u)} w(\theta) d\theta,$$

on $B_n \cap C_n$, where u is now defined by $u = \theta_0 + \frac{1}{1+\varepsilon} I^{-1}(\theta_0) \ell'_n(\theta_0)$. The right hand side of (A.14) is bounded from below by

$$(A.15) \quad \frac{m_n(X^n)}{p(X^n | \theta_0)} \geq \tilde{H}_n \left(\sum_{i=1}^3 \int_{\mathbb{R}^d} \tilde{G}_i e^{-((1+\varepsilon)n/2)(\theta-u)^t I(\theta_0)(\theta-u)} d\theta - \sum_{i=1}^3 \int_{N(\theta_0, \alpha)^c} \tilde{G}_i e^{-((1+\varepsilon)n/2)(\theta-u)^t I(\theta_0)(\theta-u)} d\theta \right).$$

Analogously to the first summation in (A.2), the summands in the first summation in (A.29) can be evaluated explicitly. They give, respectively,

$$(A.16) \quad \tilde{H}_n \tilde{G}_1 \frac{(2\pi)^{d/2}}{|n(1+\varepsilon)I(\theta_0)|^{1/2}},$$

zero, and

$$(A.17) \quad \frac{\tilde{H}_n}{2(1+\varepsilon)n} \frac{(2\pi)^{d/2} \operatorname{tr}(\nabla^2 \underline{w} I^{-1}(\theta_0))}{|n(1+\varepsilon)I(\theta_0)|^{1/2}}.$$

Next we deal with the terms in the second summation in (A.15). Note that for $\theta \in N(\theta_0, \alpha)^c$ and $\delta/(1+\varepsilon) < \alpha/2$, we have, on C_n , that $\|\theta - u\| \geq \alpha/2$. So, the first summand is bounded above by

$$(A.18) \quad \tilde{H}_n |\tilde{G}_1| \frac{(2\pi)^{d/2} 2^{d/2}}{|n(1+\varepsilon)I(\theta_0)|^{1/2}} e^{-(n/16)\alpha^2(1+\varepsilon)},$$

the second is bounded above by

$$(A.19) \quad \tilde{H}_n \frac{(2\pi)^{d/2} 2^{d/2}}{|n(1+\varepsilon)I(\theta_0)|^{1/2}} \left[\frac{2}{n(1+\varepsilon)} \tilde{G}_4^t I^{-1}(\theta_0) \tilde{G}_4 \right] e^{-(n/16)(1+\varepsilon)\alpha^2},$$

and, the third is bounded above by

$$(A.20) \quad \frac{\tilde{H}_n (2\pi)^{d/2} 2^{d/2} \sqrt{3} \sqrt{\operatorname{tr} \nabla^2 \underline{w} I^{-1}(\theta_0) \nabla^2 \underline{w} I^{-1}(\theta_0)}}{n(1+\varepsilon)|n(1+\varepsilon)I(\theta_0)|^{1/2}} e^{-(n/16)(1+\varepsilon)\alpha^2}.$$

Assembling (A.16) to (A.20) gives (2.14b). \square

PROOF OF THEOREM 2.1. First observe that for $\varepsilon \in (0, 1)$ and $\alpha, \delta > 0$ such that $\delta/(1-\varepsilon) \leq \alpha/2$ we have

$$(A.21) \quad \begin{aligned} & \frac{1}{n^{d/2}} \int w(\theta_0 | x^n) p(x^n | \theta_0) dx^n \\ & \geq \frac{w(\theta_0)}{n^{d/2}} \int_{A_n \cap B_n \cap C_n} \frac{p(x^n | \theta_0)}{m(x^n)} p(x^n | \theta_0) dx^n. \end{aligned}$$

Using (2.14a) we lower bound the right hand side of (A.21) by

$$\begin{aligned} & \frac{w(\theta_0)(1-\varepsilon)^{d/2}|I(\theta_0)|^{1/2}}{(2\pi)^{d/2}} \int_{A_n \cap B_n \cap C_n} \frac{p(x^n | \theta_0) dx^n}{H_n \left[G_1 + \frac{\operatorname{tr}[\nabla^2 \bar{w} I^{-1}(\theta_0)]}{2(1-\varepsilon)n} e^{-n(1-\varepsilon)\alpha^2/16} \right]} \\ & = \frac{(1-\varepsilon)^{d/2}|I(\theta_0)|^{1/2}}{(1+\varepsilon)(2\pi)^{d/2}} E_{\theta_0} \{ 1_{A_n \cap B_n \cap C_n} e^{-(1/2(1-\varepsilon))Z_n^t Z_n} G_5^{-1} \}, \end{aligned}$$

where $Z_n = \sqrt{n}I(\theta_0)^{-1/2}\ell'_n(\theta_0)$ and

$$(A.22) \quad G_5 = 1 + \frac{\nabla w(\theta_0)^t I^{-1/2}(\theta_0) Z_n}{\sqrt{n}(1-\varepsilon)w(\theta_0)} + \frac{Z_n^t I^{-1/2}(\theta_0) \nabla^2 \bar{w} I^{-1/2}(\theta_0) Z_n}{2(1-\varepsilon)^2 n w(\theta_0)} \\ + \frac{\text{tr} \nabla^2 \bar{w} I^{-1}(\theta_0)}{2n(1-\varepsilon)w(\theta_0)} + \frac{C_1(\theta_0) e^{-n\alpha^2(1-\varepsilon)/16}}{w(\theta_0)}.$$

By the restriction to C_n , $C_1(\theta_0)$ is bounded so the last term in the denominator in the expectation can be neglected. Since the other three nontrivial terms in the denominator sum to a small number we can apply the second order Taylor expansion of $(1+x)^{-1}$. Now the argument of the expectation is

$$(A.23) \quad 1_{A_n \cap B_n \cap C_n} \left[1 - \frac{\nabla w(\theta_0) I^{-1/2}(\theta_0) Z_n}{(1-\varepsilon)\sqrt{n}w(\theta_0)} - \frac{Z_n^t I^{-1/2}(\theta_0) \nabla^2 \bar{w} I^{-1/2}(\theta_0) Z_n}{2(1-\varepsilon)^2 n w(\theta_0)} \right. \\ \left. - \frac{\text{tr} \nabla^2 \bar{w} I^{-1}(\theta_0)}{2n(1-\varepsilon)w(\theta_0)} + \frac{(\nabla w(\theta_0)^t I^{-1/2}(\theta_0) Z_n)^2}{(1-\varepsilon)^2 n w(\theta_0)^2} \right. \\ \left. + o\left(\frac{1}{n}\right) \right] e^{-Z_n^t Z_n / (2(1-\varepsilon))}.$$

Note that by results in Clarke and Barron (1990), $P_\theta(A_n^c)$, $P_\theta(B_n^c)$ and $P_\theta(C_n^c)$ are all $o(1/n)$, so the indicator function does not affect the limiting behaviour of any of the terms in (A.23). Indeed, if one examines expression (6.3) in Clarke and Barron (1990), one sees that $P_\theta(A_n^c) = o(1/n)$ follows from including the indicator function in the second expression in (6.3) so that the left hand side of (6.3) is $o(1/n)$ and this can be used at the end of the proof of Proposition 6.3. That $P_\theta(B_n^c)$ and $P_\theta(C_n^c)$ are $o(1/n)$ follows from (4.16) and (4.17) in Clarke and Barron (1990).

Next, we obtain the limiting behavior of the six terms in (A.23). The first term in (A.23) can be written as

$$(A.24) \quad E_{\theta_0}[e^{-Z_n^t Z_n / (2(1-\varepsilon))}] - E_{\theta_0}[1_{(A_n \cap B_n \cap C_n)^c} e^{-Z_n^t Z_n / (2(1-\varepsilon))}].$$

The second term in (A.24) is $o(1/n)$. For the first term in (A.24) we use a local limit theorem from Bhattacharya and Rao (1986) (see Theorem 19.2, p. 192 and Sections 6 and 7 for definitions of quantities) so as to approximate it to order $o(1/n)$. Recall that the density f_{n,θ_0} , of Z_n , has mean zero and variance matrix equal to the identity matrix in d dimensions. Consequently, f_{n,θ_0} admits an expansion of the form

$$(A.25) \quad f_{n,\theta_0}(z) = \varphi(z) - \varphi(z) \sum_{k=1}^r \frac{p_{k,\theta_0}(z)}{n^{k/2}} + o\left(\frac{1}{n^{r/2}}\right)$$

uniformly in $z = (z_1, \dots, z_d)$, where φ is the standard normal density in d dimensions and p_{k,θ_0} is a polynomial in d arguments. We use (A.25) for $r = 2$ since we

want a $o(1/n)$ error term. The result is

$$(A.26) \ E[e^{-Z_n^t Z_n / (2(1-\varepsilon))}] = E[e^{-Z^t Z / (2(1-\varepsilon))}] - E \left[e^{-Z^t Z / (2(1-\varepsilon))} \frac{p_{1,\theta_0}(Z)}{\sqrt{n}} \right] \\ - E \left[e^{-Z^t Z / (2(1-\varepsilon))} \frac{p_{2,\theta_0}(Z)}{n} \right] + o\left(\frac{1}{n}\right),$$

where $Z \sim Normal(0, I_{d \times d})$. The first term on the right hand side of (A.26) is $(1 + \frac{1}{1-\varepsilon})^{-d/2}$ by straightforward calculations since $Z^t Z$ is χ_d^2 . The second term on the right hand side of (A.26) is zero since $p_{1,\theta_0}(Z)$ is an odd function, see Bhattacharya and Rao (1986), Expression (7.20). This leaves the third term which is difficult to evaluate in general. Expression (A.23) now gives

$$(A.27) \quad n \left[E e^{-Z_n^t Z_n / (2(1-\varepsilon))} - \left(1 + \frac{1}{1-\varepsilon}\right)^{-d/2} \right] \\ = -E e^{-Z^t Z / (2(1-\varepsilon))} p_{2,\theta_0}(Z) + o(1).$$

For the second term in (A.23) we use (A.25) with $r = 1$, since there is already a \sqrt{n} in the denominator. We have the vector equation

$$(A.28) \ E[Z_n e^{-Z_n^t Z_n / (2(1-\varepsilon))}] = E[Z e^{-Z^t Z / (2(1-\varepsilon))}] \\ - E \left[Z \frac{p_{1,\theta_0}(Z)}{\sqrt{n}} e^{-Z^t Z / (2(1-\varepsilon))} \right] + o\left(\frac{1}{\sqrt{n}}\right),$$

in which the first term of the right hand side is zero. Thus the limiting behavior of the second term in (A.23) is

$$(A.29) \quad \frac{\nabla w(\theta_0) I^{-1/2}(\theta_0)}{(1-\varepsilon)\sqrt{n}w(\theta_0)} E Z_n e^{-Z_n^t Z_n / (2(1-\varepsilon))} \\ = \frac{\nabla w(\theta_0) I^{-1/2}(\theta_0)}{n(1-\varepsilon)w(\theta_0)} E [Z p_{1,\theta_0}(Z) e^{-Z^t Z / (2(1-\varepsilon))}] + o\left(\frac{1}{n}\right).$$

The third, fourth and fifth terms are easier since they are already of order $O(1/n)$, with smaller order error terms. Thus, it is enough to use the asymptotic normality of Z_n for all of them. For the third term we obtain

$$(A.30) \quad E \left[1_{A_n \cap B_n \cap C_n} \frac{Z_n^t I^{-1/2}(\theta_0) \nabla^2 \bar{w} I^{-1/2}(\theta_0) Z_n e^{-Z_n^t Z_n / (2(1-\varepsilon))}}{2(1-\varepsilon)^2 w(\theta_0)} \right] \\ \rightarrow \frac{E [Z^t I^{-1/2}(\theta_0) \nabla^2 \bar{w} I^{-1/2}(\theta_0) Z e^{-Z^t Z / (2(1-\varepsilon))}]}{2(1-\varepsilon)^2 w(\theta_0)} \\ = \frac{\text{tr}[\nabla^2 \bar{w} I^{-1}(\theta_0)]}{2(1-\varepsilon)^2 w(\theta_0) \left(1 + \frac{1}{1-\varepsilon}\right)^{d/2+1}};$$

for the fourth term we obtain

$$(A.31) \quad \frac{\text{tr} \nabla^2 w(\theta_0) I^{-1}(\theta_0)}{2(1-\varepsilon)w(\theta_0)} E_{\theta_0} [1_{A_n \cap B_n \cap C_n} e^{-Z_n^t Z_n / (2(1-\varepsilon))}] \\ \rightarrow \frac{\text{tr} \nabla^2 w(\theta_0) I^{-1}(\theta_0)}{2(1-\varepsilon)w(\theta_0) \left(1 + \frac{1}{1-\varepsilon}\right)^{d/2}};$$

and for the fifth term we obtain

$$(A.32) \quad E \left[1_{A_n \cap B_n \cap C_n} e^{-Z_n^t Z_n / (2(1-\varepsilon))} \frac{[\nabla w(\theta_0)^t I^{-1/2}(\theta_0) Z_n]^2}{(1-\varepsilon)^2 w(\theta_0)^2} \right] \\ \rightarrow \frac{\nabla w(\theta_0)^t I^{-1/2}(\theta_0) \nabla w(\theta_0)}{(1-\varepsilon)^2 w(\theta_0)^2 \left(1 + \frac{1}{1-\varepsilon}\right)^{d/2+1}}.$$

Putting together (A.27), (A.29) through (A.32) and letting n increase gives the stated lower bound.

The upper bound follows by writing the expected posterior as a sum of two integrals, one over $B_n \cap C_n$ which can be controlled as in the lower bound, and one over $(B_n \cap C_n)^c$ which gives the term involving J_n . \square

Appendix B: Proofs from Section 3

Here we give the proof of Theorem 3.1. We will only prove the lower bound (3.2a). The proof of (3.2b) is similar. First observe that for $\varepsilon \in (0, 1)$ and $\alpha, \delta > 0$ such that $\delta/(1-\varepsilon) \leq \alpha/2$ we have

$$(B.1) \quad \frac{1}{n^d} \int w^2(\theta_0 | x^n) p(x^n | \theta_0) dx^n \\ \geq \frac{w^2(\theta_0)}{n^d} \int_{A_n \cap B_n \cap C_n} \left\{ \frac{p(x^n | \theta_0)}{m(x^n)} \right\}^2 p(x^n | \theta_0) dx^n.$$

Using (2.14a) we lower bound the right hand side of (B.1) by

$$\frac{w^2(\theta_0)(1-\varepsilon)^d |I(\theta_0)|}{(2\pi)^d} \\ \int_{A_n \cap B_n \cap C_n} \frac{p(x^n | \theta_0) dx^n}{H_n^2 \left[G_1 + \frac{\text{tr}[\nabla^2 \bar{w} I^{-1}(\theta_0)]}{2(1-\varepsilon)n} + C_1(\theta_0) e^{-n(1-\varepsilon)\alpha^2/16} \right]^2} \\ = \frac{(1-\varepsilon)^d |I(\theta_0)|}{(1+\varepsilon)(2\pi)^d} E_{\theta_0} \{ 1_{A_n \cap B_n \cap C_n} G_5^{-2} e^{-Z_n^t Z_n / (1-\varepsilon)} \},$$

where $Z_n = \sqrt{n} I(\theta_0)^{-1/2} \ell'_n(\theta_0)$ and G_5 is given by (A.22). By the restriction to C_n , $C_1(\theta_0)$ is bounded so the last term in the denominator in the expectation can

be neglected. Since the other three nontrivial terms in the denominator sum to a small number we can apply the second order Taylor expansion of $(1+x)^{-2}$. Now the argument of the expectation is

$$(B.2) \quad 1_{A_n \cap B_n \cap C_n} \left\{ 1 - \frac{2\nabla w(\theta_0)I^{-1/2}(\theta_0)Z_n}{(1-\varepsilon)\sqrt{nw}(\theta_0)} - \frac{Z_n^t I^{-1/2}(\theta_0)\nabla^2 \bar{w} I^{-1/2}(\theta_0)Z_n}{(1-\varepsilon)^2 nw(\theta_0)} \right. \\ \left. - \frac{\text{tr} \nabla^2 \bar{w} I^{-1}(\theta_0)}{n(1-\varepsilon)w(\theta_0)} + \frac{3(\nabla w(\theta_0)^t I^{-1/2}(\theta_0)Z_n)^2}{(1-\varepsilon)^2 nw(\theta_0)^2} + o\left(\frac{1}{n}\right) \right\} e^{-Z_n^t Z_n / (1-\varepsilon)}.$$

As before, $P_\theta(A_n^c)$, $P_\theta(B_n^c)$ and $P_\theta(C_n^c)$ are all $o(1/n)$ so the indicator function does not affect the limiting behaviour of any of the terms in (B.2). Let us consider the limiting behavior of the six terms in (B.2).

The first term in (B.2) can be written as

$$(B.3) \quad E_{\theta_0}[e^{-Z_n^t Z_n / (1-\varepsilon)}] - E_{\theta_0}[1_{(A_n \cap B_n \cap C_n)^c} e^{-Z_n^t Z_n / (1-\varepsilon)}].$$

The second term in (B.3) is $o(1/n)$. For the first term in (B.3), we use the expansion (A.24) with $r = 2$ for density f_{n,θ_0} , of Z_n . The result is

$$(B.4) \quad E[e^{-Z_n Z_n / (1-\varepsilon)}] = E[e^{-Z^t Z / (1-\varepsilon)}] - E\left[e^{-Z^t Z / (1-\varepsilon)} \frac{p_{1,\theta_0}(Z)}{\sqrt{n}}\right] \\ - E\left[e^{-Z^t Z / (1-\varepsilon)} \frac{p_{2,\theta_0}(Z)}{n}\right] + o\left(\frac{1}{n}\right),$$

where $Z \sim \text{Normal}(0, I_{d \times d})$. The first term on the right hand side of (B.3) is $(1 + \frac{2}{1-\varepsilon})^{-d/2}$ by straightforward calculations since $Z^t Z$ is χ_d^2 . The second term on the right hand side of (B.3) is zero since $p_{1,\theta_0}(Z)$ is an odd function, see Bhattacharya and Rao (1986), Expression (7.20). This leaves the third term which is difficult to evaluate in general. Expression (B.3) now gives

$$(B.5) \quad n \left[E e^{-Z_n Z_n / (1-\varepsilon)} - \left(1 + \frac{2}{1-\varepsilon}\right)^{-d/2} \right] = -E[e^{-Z^t Z / (1-\varepsilon)} p_{2,\theta_0}(Z)] + o(1).$$

For the second term in (B.2) we use (A.24) with $r = 1$, since there is already a \sqrt{n} in the denominator. We have the vector equation

$$(B.6) \quad E[Z_n e^{-Z_n^t Z_n / (1-\varepsilon)}] = E[Z e^{-Z^t Z / (1-\varepsilon)}] \\ - E\left[Z \frac{p_{1,\theta_0}(Z)}{\sqrt{n}} e^{-Z^t Z / (1-\varepsilon)}\right] + o\left(\frac{1}{\sqrt{n}}\right)$$

in which the first term of the right hand side is zero. Thus the limiting behavior of the second term in (B.2) is

$$(B.7) \quad \frac{2\nabla w(\theta_0)I^{-1/2}(\theta_0)}{(1-\varepsilon)\sqrt{nw}(\theta_0)} E[Z_n e^{-Z_n^t Z_n / (1-\varepsilon)}] \\ = \frac{2\nabla w(\theta_0)I^{-1/2}(\theta_0)}{n(1-\varepsilon)w(\theta_0)} E[Z p_{1,\theta_0}(Z) e^{-Z^t Z / (1-\varepsilon)}] + o\left(\frac{1}{n}\right).$$

The third, fourth and fifth terms are easier since they are already of order $O(1/n)$, with smaller order error terms. Thus, it is enough to use the asymptotic normality of Z_n for all of them. For the third term we obtain

$$\begin{aligned}
 \text{(B.8)} \quad E & \left[1_{A_n \cap B_n \cap C_n} \frac{Z_n^t I^{-1/2}(\theta_0) \nabla^2 \bar{w} I^{-1/2}(\theta_0) Z_n e^{-Z_n^t Z_n / (1-\varepsilon)}}{(1-\varepsilon)^2 w(\theta_0)} \right] \\
 & \rightarrow \frac{1}{(1-\varepsilon)^2 w(\theta_0)} E[Z^t I^{-1/2}(\theta_0) \nabla^2 \bar{w} I^{-1/2}(\theta_0) Z e^{-Z^t Z / (1-\varepsilon)}] \\
 & = \frac{\text{tr} \nabla^2 \bar{w} I^{-1}(\theta_0)}{(1-\varepsilon)^2 w(\theta_0) \left(1 + \frac{2}{1-\varepsilon}\right)^{d/2+1}};
 \end{aligned}$$

for the fourth term we obtain

$$\begin{aligned}
 \text{(B.9)} \quad \frac{\text{tr} \nabla^2 w(\theta_0) I^{-1}(\theta_0)}{(1-\varepsilon) w(\theta_0)} E_{\theta_0} [1_{A_n \cap B_n \cap C_n} e^{-Z_n^t Z_n / (1-\varepsilon)}] \\
 \rightarrow \frac{\text{tr} \nabla^2 w(\theta_0) I^{-1}(\theta_0)}{(1-\varepsilon) w(\theta_0) \left(1 + \frac{2}{1-\varepsilon}\right)^{d/2}};
 \end{aligned}$$

and for the fifth term we obtain

$$\begin{aligned}
 \text{(B.10)} \quad E & \left[1_{A_n \cap B_n \cap C_n} e^{-Z_n^t Z_n / (1-\varepsilon)} \frac{3[\nabla w(\theta_0)^t I^{-1/2}(\theta_0) Z_n]^2}{(1-\varepsilon)^2 w(\theta_0)^2} \right] \\
 & \rightarrow \frac{3 \nabla w(\theta_0)^t I^{-1/2}(\theta_0) \nabla w(\theta_0)}{(1-\varepsilon)^2 w(\theta_0)^2 \left(1 + \frac{2}{1-\varepsilon}\right)^{d/2+1}}.
 \end{aligned}$$

Putting together (B.5) through (B.10) gives the stated lower bound.

REFERENCES

Bernardo, J. M. (1979). Reference posterior distributions for Bayesian inference, *J. Roy. Statist. Soc. Ser. B*, **41**, 113–147.

Bhattacharya, R. N. and Rao, R. R. (1986). *Normal Approximation and Asymptotic Expansions*, Krieger Publishing, Malabar, Florida.

Casella, G. and Berger, R. L. (1990). *Statistical Inference*, Wadsworth and Brooks/Cole, Pacific Grove, California.

Clarke, B. and Barron, A. R. (1990). Information-theoretic asymptotics of Bayes methods, *IEEE Trans. Inform. Theory*, **36**(3), 453–471.

Clarke, B. and Sun, D. (1997). Reference priors under the Chi-square distance, *Sankhyā Ser. A*, **59**, 215–231.

De Bruijn, N. G. (1961). *Asymptotic Methods in Analysis*, 2nd ed., Dover Publications, New York.

Feller, W. (1971). *An Introduction to Probability Theory and Its Applications*, Vol. II, 2nd ed., Wiley, New York.

Kass, R. E. and Raftery, A. E. (1995). Bayes factors, *J. Amer. Statist. Assoc.*, **90**, 773–795.